## CASEVILLE, MICH., JUNE 29, 1894. BY DR.J.T.SCOVELL.

Thoken kund

There is no gill net fishing from Caseville; but there are several strings of pound nets along the shore near Caseville.

P

(1). An old man by the name of Germain has a string of 1 old net extending north-westerly from Oak Point, about 2 1/2 miles northerly from Caseville. The leader is about 20 rods long, water 4 feet and the pot 20 x 20 x 12 standing in 12 feet of water. The pot is just in and no fish have been marketed.

P.

(2). About one mile below or northerly from Caseville, F.Marsac and Eugene Willett have a string of 4 pounds, commencing 50 rods from shore in 4 feet of water.

1st leader 50 rods, hearts 6 rods, pot 20

P

x 20 x 10, 10 feet of water.

2d leader 75 rods, hearts 6 rods, pot 20 x 20 x 13, 13 feet of water.

3d leader, 75 rods, hearts 6 rods, pot 20 x 20 x 15, 15 feet of water.

4th leader 75 rods, hearts 6 rods, pot 20 x 20 x 18, 18 feet of water.

The nets set about north-west. They are of cotton twine No.12. Mesh 6 1/2-5-and 2 1/2, 2-inch back. The twine shrinks but little. The nets were torn out and much damaged by the storm of May 20, but such storms are not common and usually not much damage is done. The bottom is sandy, with many scattered boulders and some bed rock, so that it is somewhat difficult to find grounds on which stakes can be driven. catch is mainly pickerel, herring, whitefish, sturgeon, some trout. Very few black bass, about 130 pounds this season.

Shows

Collin

Catch

Collecting Seiner

Seined in Pigeon River above the town of Caseville where it was a series of rapids and quiet pools; averaging parhaps 15 feet wide and 4 inches deep. Found several species of fish and a bivalve shell fish, which looked like the shells we found in the stomach of the whitefish. Fished in some bayous along the shore, also in the shallow waters of the lake--several species.

(3). ALEXANDER TRUDELL. Has a string of 2 pounds set about 2 miles above the town of Caseville. The leader begins in about 6 feet of water about 50 rods from shore and extends north-westerly.

1st leader 100 rods, 7, 30 x 30 x 14 in 13 feet of water.

2d leader 80 rods, 7, 30 x 30 x 15 in 16 feet of water.

The second pot has not been out this sea-

Walleys Ourale.

son. Cotton twine No.12. Mesh 6-4-3 with 2 1/2 back. The bottom of clay and sand with many scattering boulders. Catch pickerel, those smaller than 1/2 pound thrown back into bay. Also catch catfish, sturgeon, saugers, perch, mullet, suckers, sheepshead. Saw the above taken from the net. Whitefish and trout sometimes taken and an occasional b.bass. From the dock south-westerly a bar of sand and gravel. Seems to be a good spawning ground for b.bass in June. During July and August no nets out. Set again in September and catch herring, pickerel, white-

Shauming!

Scasin

catch

when lighted

Ship fishing

Lift every morning if convenient.

fish, lake perch, mullet and suckers.

Thinks fish would be about as abundant as ever if there were no fishing for five years.

(4). JAMES REEVES. About 1 1/2 or 2 miles above Trudell's nets. Mr.Reeves has

P.

T?

been in the business about 14 years. Has a string of 8 pounds. They commence in 6 feet of water about 80 rods from shore and extend about north-west.

lst leader 6 rods, heart 5 rods, pot 28
x 28 x 12, 12 feet of water.

2d leader 60 rods, heart 5 rods, pot 30 x 30 x 14, 14 feet of water.

3d leader 60 rods, heart 5 rods, pot 30 x 30 x 16 in 16 feet of water.

4th leader 60 rods, heart 5 rods, pot 30 x 30 x 18 in 18 feet of water.

5th leader 60 rods, heart 5 rods, pot 30 x 30 x 19 in 19 feet of water.

6th leader 60 rods, heart 5 rods, pot 30 x 30 x 20 in 20 feet of water.

7th leader 60 rods, heart 5 rods, pot 30 x 30 x 21 in 21 feet of water.

8th leader 60 rods, heart 5 rods, pot 30 x 30 x 22 in 22 feet of water.

F.

The nets are made of No.12 and 15 cotton twine, which shrinks a little.

Mesh 5-4-2 1/2 inches, 2-inch back.

Season

Nets put out about April 15, taken out about July 5. The catch mainly of herring, pickerel, whitefish, sturgeon, perch, suckers, mullet, trout, catfish, sheepshead, etc.

cut off hout

Thinks gill nets at mouth of bay cuts off the trout. The herring and whitefish more abundant in the fall season, which begins about September 15.

Any fish caught weighing one pound or

Small fish

less, if in good condition, are thrown back into the bay. Early in the season the inside nets seem to do the best; later the outside ones are best. The fish going into deeper water. During the fall season all

June outup.

One string of pound nets does interfere

the pounds fish about the same, depending

much on the weather.

The string with

with others, especially when the fish are moving up or down the coast, as the nets are usually set perpendicular to the coast so as to eatch the fish moving in that way. The pickerel spawn in the spring on sand or gravel in shallow, warm water. Only catches a few b.bass, not more than 200 pounds this season. Bottom of his ground sand or elay, no boulders. Was much damaged by the storm of May 20, 1894, but damage from storms is

(5). LOUIS TRUDELL, of Bay City, Mich.

Has a string of 9 pounds situated about 2

miles S.W. of Little Charities Island. The

string stands about N.W.

usually very slight.

lst leader 80 rods, hearts 5 rods, pot 32 x 32 x 16 in 16 feet of water.

2d leader 80 rods, hearts 5 rods, pot 32 x 32 x 18 in 18 feet of water.

nalleys Spanning

bussi

Lours

P

3d leader 80 rods, hearts 5 rods, pot 32 x 32 x 20 in 20 feet of water.

And so on, the last pot standing in 32

feet of water. Nets made of cotton twine No. 15 thread. Mesh 6-5-2 1/2 and 2-inch back. Bottom of clay, good driving. Outside nets most profitable. Formerly fishing was better in shoel water, now it is better in deep water. In stormy weather fish go into deep water. Some localities that seem fished out, if abandoned, become good again after 4 or 5 years' rest. Spring catch 1/2 pickerel, 1/2 whitefish. The pickerel are caught as they are going out of the bay from their spawning grounds, which is in streams and shoal water. Catch also perch, herring, sturgson, trout

Catch

Fall catch 2/3 whitefish, 1/3 pickerel, herring and some trout. Catch spawning

and others. No b.bass. Never catch spawn-

catch fall willing willing

ing pickerel.

P. Williaming, with

Qual meth

foulwarder

market

aut it

will local

Herring is

-0

whitefish in the last of November. Money is mainly made on the herring. Fishing would be as good as ever if the small or young fish were not caught in small mesh nets.

Small mesh fishing in the rivers bad. 12 years ago 3 pounds were as profitable as 9 pounds are now. The foul water of the river hurts river fishing, and fishing in the head of the bay. The catch is brought to Bay City. The herring salted, the others sold fresh.

Spring catch 8 pound nets. . . \$1,300
Fall \* 9 \* \* . . . 4,500

Whitefish found on sandy, gravelly or stony bottom. They eat mollusks, water bugs, May flies and perhaps eat perch spawn, and some algea, herring much like whitefish.

(6). R.L.GILLINGHAM, living in Bay Port, Mich. Has a string of 7 nets, commencing in P

14 feet of water about 2 miles north of Sand
Point or Point Charities and about 2 miles
westerly from J.Reeves' nets and extending
north-east.

The string begins with a pot 30 x 30 x 16 in 14 feet of water on a sand and clay bot-tem.

1st leader 80 rods, 6 rods, 30 x 30 x 18 in 16 feet of water.

2d leader 80 rods, 6 rods, 30 x 30 x 19 in 17 feet of water.

3d leader 80 rods, 6 rods, 30 x 30 x 20 in 18 feet of water.

4th leader 80 rods, 6 rods, 30 x 30 x 20 in 18 feet of water.

5th leader 80 rods, 6 rods, 30 x 30 x 20 in 18 feet of water.

6th leader 80 rods, 6 rods, 30 x 30 x 20 in 18 feet of water.

Bottom of outside nets sand covering a

bollin

P.

seasin

calch

mil.

bed of clay. No boulders. The direction is about perpendicular with bar from Sand Point. Mesh 6-4-3. Pot 2 1/2 and 2 back. Cotton twine No.12 or 15. Sets out as soon as possible after the opening of navigation and pulls out after July 5 to 10. Spring catch pickerel, herring, perch, whitefish, trout, catfish, sturgeon for market. Also suckers, mmallet and sheepsheed. Fall catch herring, whitefish, pickerel, perch, suckers and mullet, sheepshead, etc. Mr.Gillingham says that this spring while raising pots he has seen great numbers of small whitefish escaping. These he thinks are those that have

## BAY PORT, MICH.

Lescrip

been planted.

Situated on Wild Fowl Bay, a shallow sheet of water; shut in by islands, so as to be nearly land locked, the bottom is general-

T. by the

ly sandy and muddy, great areas being covered with bullrushes and different kinds of sedges and grasses. There are several strings of nets outside the islands in the main bay and many little fyke nets or gobblers and some pounds in the little bay. The town of Bay Port is near the center of Sec. 36, T.9 E., R.17 north. North Island is mainly in Sec. 21, same town. and range. Commencing about the center of Sec. 21, 80 rods from shore in 8 feet of water is a string of 22 pounds extending north-west by west into the main bay.

The nets are owned by J.& J.Gillingham and W.C.Gillingham, of Bay Port, Mich. These nets are about 5 miles south of R.L.Gillingham's nets.

lst leader 80 rods, hearts 6 rods, pot 30 x 30 x 17 in 15 feet of water.

2d leader 80 rods, hearts 6 rods, pot 30

P.

x 30 x 20 in 18 feet of water.

3d leader 80 rods, hearts 6 rods, pot 30

x 30 x 20 in 18 feet of water.

4th leader 80 rods, hearts 6 rods, pot 30

x 30 x 20 in 18 feet of water.

5th leader 80 rods, hearts 6 rods, pot 30

x 30 x 21 in 19 feet of water.

In this leader there is a gap.

6th leader 80 rods, hearts 6 rods, pot 30

x 30 x 21 in 19 feet of water.

7th leader 80 rods, hearts 6 rods, pot 30

x 30 x 18 in 16 feet of water.

There a gap.

And pound or pot No.3 30 x 30 x 15 in 14 feet of water.

8th leader 80 rods, hearts 6 rods, pot 30

x 30 x 15 in 13 feet of water.

9th leader 80 rods, hearts 6 rods, pot 30

x 30 x 14 in 12 feet of water.

10th leader 80 rods, hearts 6 rods, pot

P

 $30 \times 30 \times 12$  in 10 feet of water.

11th leader 80 rods, hearts 6 rods, pot

30 x 30 x 12 in 10 feet of water.

12th leader 80 rods, hearts 6 fods, pot

30 x 30 x 12 in 10 feet of water.

There a gap and pot 14 30 x 30 x 13 in 11 feet of water.

13th leader 80 rods, hearts 6 rods, pot 30 x 30 x 13 in 11 feet of water.

There a gap and pot  $16 30 \times 30 \times 13$  in 11 feet of water.

14th leader 80 rods, hearts 6 rods, pot

30 x 30 x 14 in 12 feet of water.

15th leader 80 rods, hearts 6 rods, pot

30 x 30 x 15 in 13 feet of water.

16th leader 80' rods, hearts 6 rods, pot

 $30 \times 30 \times 15$  in 13 feet of water.

17th leader 80 rods, hearts 6 rods, pot

 $30 \times 30 \times 15$  in 13 feet of water.

18th leader 80 rods, hearts 6 rods, pot

 $30 \times 30 \times 17$  in 15 feet of water.

19th leader 80 rods, hearts 6 rods, pot

P.

 $30 \times 30 \times 18$  in 16 feet of water.

Extending nearly 6 miles into the main bay. These nets are made of No.12, 15 and 16 cotton twine and are 2 feet deeper than the water. The mesh is 6-4-2 1/2 sides and 2-inch back. Somewhat larger at first, but shrink. The nets are put out as soon as navigstion opens in the spring and are taken out about July 1. Fall fishing begins about September 10 and lasts till about November 15, according to season. Spring catch pickerel, herring, perch, some whitefish, sturgeon, also some suckers, mullet, sheepsheads, &c. Fall catch herring, whitefish, pickerel, perch, suckers. Storms injure the nets somewhat, especially S.W. and N.E.gales, but the loss is usually not great. Nets are sometimes injured by vessels running through them, but the damage is usually light. Be-

fore 1870 fishing was good, them for 5 years

Calch

Storms

Justing history

Slurgen

but few fish were caught, but in 1878 and 1879 began again and now are catching more fish than before. Sturgeon were formerly very, very abundant, but now only a few are caught each year, and the catch is gradually

bass

decreasing. Only a very few b.bass are caught in the pound nets. When caught, they must be taken out soon or they may escape. Trout, pickerel, bass and similar fish escape from nets more readily and in greater numbers than whitefish and herring. Have

herring

never seen spawning pickers! on these grounds, think all have spawned before reach-

herring Shewar

ing them. Have seen a few spawning whitefish and herring, but usually we have pulled out

Smalefin

pound thrown away. All fish of 1 pound and

before they begin to spawn. Fish below 3/4

bollim' a sand, clay and cobble stone. Herring gill

extensively in the pots when they are lifted,

gilling Herring walley

and sometimes pickerel gill in the pounds and points of the hearts. Some catfish are caught with the hook during the summer. Some ice fishing is done with the spear, some 12 to 15 shanties about these islands. They catch grass pike, perch and black bass. Market fish mainly in Bay City, Detroit and New York. Some sold to peddlers and in land towns. The herring are salted and sold in Detroit. Other fish sold fresh. Ten men depend entirely on Tishing. 25 men fish from 3 to 5 months each year, and a few, perhaps 10 more, work only in heavy herring fishing in the fall. This information is mainly from W.C.Gillingham, of Gillingham Bros. who are among the best fishermen on the bay.

The Gillingham Bros. and others are of the opinion that whitefish planted by the State and U.S. Commissions are now being caught along the west shores of Saginaw Bay. Fish

offal is generally buried. Sometimes thrown into the bay. Offal hurt the grounds for trout and whitefish, but not for pickerel and perch. Logs floating about the bay bother fishermen. Mills along the river and bay furnish material that fouls the water of the bay. Water unusually bad this spring and some lay it to the pumping of mine water

into the bay at Sebewaing. W.C.Gillingham

has 3 boats, R.L.Gillingham has 2 and J.& J.

Logs Sawmill

Gillingham 6.

LOUIS DUBEY, Bay Port, Mich. fished on these grounds some 9 years. He has a string of 17 pounds about N.W.by west, commencing about 80 rods from the shore of Heisterman's Island, near the south-west corner of Sec. 29, T.9 B., R.17 N., in about 13 feet of water.

1st leader 90 rods, hearts 5 rods, pot 30

P

x 30 x 13 in 13 feet of water.

2d leader 80 rods, hearts 5 rods, pot 30 x 30 x 14 in 14 feet of water.

3d leader 80 rods, hearts 5 rods, pot 30 x 30 x 17 in 17 feet of water.

4th leader 70 rods, hearts 5 rods, pot 30 x 30 x 18 in 18 feet of water.

5th leader 70 rods, hearts 5 rods, pot 30 x 30 x 19 in 19 feet of water.

6th leader 70 rods, hearts 5 rods, pot 30 x 30 x 18 in 18 feet of water.

7th leader, there a gap of 20 rods and pot 30  $\times$  30  $\times$  17 in 17 feet of water.

8th leader, 75 rods, hearts 5 rods, pot 30 x 30 x 11 in 11 feet of water.

9th leader 75 rods, hearts 5 rods, pot 30 x 30 x 10 in 10 feet of water.

10th leader 75 rods, hearts 5 rods, pot 30 x 30 x 9 in 9 feet of water.

11th leader 75 rods, hearts 5 rods, pot

P.

P.

30 x 30 x 9 in 9 feet of water.

12th leader 75 rods, hearts 5 rods, pot 30 x 30 x 9 in 9 feet of water.

The above 12 fished by Dubey himself.

Beyond these in the same string are 4 leaders and 5 pots, all in about 12 feet of water, fished by Peter Whitman.

13 pot 30 x 30 x 12 in 12 feet of water.

14th leader 75 rods, hearts 5 rods, pot

 $30 \times 30 \times 12$  in 12 feet of water.

15th leader 75 rods, hearts 5 rods, pot 30 x 30 x 12 in 12 feet of water.

16th leader 75 rods, hearts 5 rods, pot 30 x 30 x 12 in 12 feet of water.

17th leader 75 rods, hearts 5 rods, pot  $30 \times 30 \times 12$  in 12 feet of water.

The mesh is 6 in.,4 1/2 and 5, 2 1/2 sides and 2-inch back. Small mesh is necessary, as much of the catch is herring, only a few small pickerel caught. Not more than

Catch herring wally aut, eatch anall walley

catch

1,000 pounds, weighing 1 pound or less this season. Bottom of the ground sand and gravel, some mud with clay beneath. Catch pickerel, herring, trout, whitefish, sturgeon, grass pike, muskallonge, rock bass, black bass, perch, suckers, mullet, etc., etc.

movement

si umming

beat spring

Variation in Bay princing

Storms.

millest pering fall catch herring

Pickerel go up the bay to spawn and we catch them on their way out or back so that we never catch spawning pickerel. And we never catch spawning whitefish, as we have to take out nets before they or the herring spawn. Spring catch best is May and June. On account of storms and shallow water, there is more variations in bay fishing than in lake fishing. The weather seems to be getting more and more stormy and more damage is done by storms, ice, etc., than formerly. Spring catch mainly pickerel, some herring. Fall catch mainly herring, a few others. The herring seem to be holding their own,

wally decrease

but the pickerel perhaps are falling off.

Whitefish seem to be coming back as the water rests; but not from planting. Fish go out as water falls and come back as it rises.

foulwater

Moving ice drives fish away. Water of the bay getting too foul for good fishing. Shallow water gets foul under the ice and fish are driven away. We send the pickerel to Bay City without dressing. The herring are

quarket

salted.

P. Inhus

CHARLES VANDEMARK. Has a string of 7 gobblers and pound nets south of Island in Sec.17, T.9 east, Range 16 north, commencing in 4 feet of water about 30 rods from shore and extending about S.W.

1st leader 20 rods, hearts 2 rods, gobbler 16 feet, 5 to 6 feet of water.

2d leader 16 rods, hearts 2 rods, gobbler 16 feet, 5 to 6 feet of water.

3d leader 13 rods, hearts 2 rods, gobbler 16 feet, 5 1/2 feet of water.

4th leader 19 rods, hearts 2 rods, gobbler 16 feet, 5 1/2 feet of water.

5th leader 30 rods, hearts 5 rods, pot 20 x 20 x 9 in 9 feet of water.

6th leader 30 rods, hearts 5 rods, pot 20 x 20 x 9 in 9 feet of water.

7th leader 20 rods, hearts 2 rods, gobbler 16 feet in 7 feet of water.

Calch

The catch is mainly perch, suckers, pickerel and grass pike. Mesh 5-4-2 1/2 and 2-

collin

ek og skrib i

depth

inch back. A few black bass are caught.

Bottom sandy and muddy. This is practically shallow water fishing, but the nets extend across a channel, and it is claimed that they prevent fish from coming into the bay and lessen the catch of the inside fishermen.

In the N.E. corner of Wild Fowl Bay on the

S.W. Sec. 8, T.10 east, Range 17 N. >

tykes

JOHN ROW AND HARRY SEIPS. Have six fyke nets or gobblers. The leaders are about 20 rods each, hearts 38 feet and the gobblers 20 feet. Standing on a mud bottom in about 4 feet of water east and west. Mesh originally 5-3-and 2 1/2 inches, but the twine has shrunken so that some of the mesh is less than 2 inches. Somewhat sheltered by banks of rushes so that there is little damage from storms. Catch mainly catfish, perch, black bass, rock bass and suckers. Mr. Row could give no information about fish or capacity of nets, as he had only been fishing about 3 months.

S forms

calin

wm.ROW AND CHARLES WYMAN. Are fishing a string of 3 pounds near the mouth of Mud Creek on Sec. 17, T.10 E., R.17 north. Com-

mence about 1/2 mils from shore in 2 1/2 feet of water extending westerly or south of west.

lst leader 50 rods, hearts 3 rods, pot 30 x 30 x 8 in 8 feet of water.

2d leader 40 rods, hearts 3 rods, pot 30 x 30 x 9 in 9 feet of water.

3d leader 45 rods, hearts 3 rods, pot 30 x 30 x 11 in 11 feet of water.

Made of cotton No.12 thread. Mesh 5-3and 2 1/2, when measured it was scant 2-inch. Catch, perch, catfish, mullet, suckers, pickerel, rock bass, some black bass and others. Mud bottom.

Catch

Collin

fyties,

Standing north-westerly from the northeast corner of Sec.36, T.9-16, there is a string of 6 fyke nets or gobblers which are fished by Eli Britt and Lobdale. Beginning in 2 1/2 feet of water about 3/4 of a mile from shore. fyker.

1st leader 12 rods, hearts 18 feet, gobbler 13 feet, 2 1/2 feet of water.

2d leader 6 rods, hearts 18 feet, gobbler 13 feet, 3 1/2 feet of water.

3d leader 7 rods, hearts 18 feet, gobbler 13 feet, 4 1/2 feet of water.

And so on. Cotton No.12 twins. Mesh 5 1/2-4-and 3 inches. Clay or sandy bottom with many boulders. Catch mainly perch, pickerel, pike, rock bass, suckers, catfish and an occasional black bass.

# yher

Due west of Bay Port on the west line of Sec.35-9-17, Frank Leveno has 3 gobblers with about 6 rod leaders, 15 ft.hearts and 15 ft.gobblers. Cotton No.12 twine with lawful mesh 5-4-and 3 1/2 inches. Catching mainly black bass, only occasionally suckers, perch, &c. About 1/2 mile S.W.he has another similar to the ones described. Muddy bottom.

Calch

fyker bottom Jykes.

GEORGE MILLER, on Heisterman's Island, on Sec. 28, T.9 E., R.17 north. Fishes a string of from 3 to 6 gobblers. The leaders are about 10 rods long, hearts about 20 feet and gobblers 12 feet. Mesh 4-3 1/2-2 1/2 to 3 for bag. Catch perch, grass pike, suckers, rock bass, some black bass and some catfish, mullet, &c. Bottom sandy or muddy. Sets in different places and directions in from 4 to 6 feet of water. Fishes spring and fall. not much in winter or summer. Fish go out in summer on account of warm water. Go out in winter on account of shallow water getting foul under ice. More fish in high water, less in low. Think outside fishermen should have larger mesh twine, which would be better for the fishing industry. The av-

Calch

Collin

loc.

Scalu

fort and ment

should use hugh large in outside

ant, calch

in 24 hours.

West of the mouth of the Shebeon Creek,

erage catch for a gobbler is about 25 pounds

Lypun

near the N.W.corner of Sec.15, T.9 E.,R.16
W.,Lafayette Evans has stakes out for 3 gobblers, with 10 rod leaders, 25 ft.hearts and
16 pots: Standing on muddy bottom in 4 feet

bollim

of water. When fishing he catches perch,

calet

suckers, bullheads, catfish, rock bass,

black bass, pickerel, pike and others. Mesh

4-3 1/2-2 1/2 inches. Nets stend nearly

4-3 1/2-2 1/2 inches. Nets stand nearly

Just movements

come in with water. The outside nets cut

east and west. Fish go out with water and

out fish and lessen the catch of the inside

twine.

in the same neighborhood, but are not in the

\_

L.H.FITTENGER. Has 5 gobblers which he sometimes fishes on the N.W. Sec.16, T.9 E., R.16 N. The leaders are about 10 rods each, hearts about 20 feet and nets about 16 feet.

Jeykes.

He generally sets the nets north and south in about from 3 to 5 feet of water. Mesh 5-4 and scant 2 inches after shrinkage.

Catch catfish, perch, suckers and a few bass.

On Sec. 21, 9 E., R.16 N., Henry Quick has two small pounds. Commences near the shore in 2 feet of water.

1st leader 40 rods, hearts 60 feet, pot 20 x 24 x 8 in 6 feet of water.

2d leader 40 rods, hearts 60 feet, pot 20 x 24 x 8 in 5 feet of water.

Ground gravel, clay and stone. Wesh 5-

5-3 with 2-inch back. Think it would be bet-

ter for the fishermen to use larger mesh.

fish, pickerel, herring, sturgeon and cat-

Catch black bass, pike, perch, mullet, bill-

in Shebeon River, but now they are seldom

fish. Says muskallonge used to be abundant

seen,

On Sec. 28-9-B., 16 N., Mr. David Girard has

catch

: h. ---

Shouldre se Larjer mush

catch

maskin

Types

catel

to 15 rods. Hearts from 20 to 25 feet and the gobblers about 16 feet each. They stand a little north of west in about 4 feet of water on a sandy or maddy bottom. He sets out his note early in May and takes them up early in July, putting out again in October. Mesh 5-3 1/2-2 1/2 inches. Catch perch. pike, dogfish, bullheads, black bass, etc. Perch yield best just after their spawning season. The nets are about 1 mile south of Henry Quick's nets. About 4 miles northwest of the town of Sebewaing, John Pitcher has a string of 2 pounds. Standing nearly east and west in about 12 feet of water.

Mesh 5-4-2 1/2 inches. Catch pickerel and

herring. Only this spring on this ground.

a string of six gobblers. Leaders from 10

cotch